arXiv:1510.01283v2 [math.AT] 29 Oct 2015

THE η-INVERTED $\mathbb{R}$-MOTIVIC SPHERE

BERTRAND J. GUILLOU AND DANIEL C. ISAKSEN

Abstract. We use an Adams spectral sequence to calculate the $\mathbb{R}$-motivic stable homotopy groups after inverting η. The first step is to apply a Bockstein spectral sequence in order to obtain h_1-inverted $\mathbb{R}$-motivic Ext groups, which serve as the input to the η-inverted $\mathbb{R}$-motivic Adams spectral sequence. The second step is to analyze Adams differentials. The final answer is that the Milnor-Witt $(4k-1)$-stem has order 2^{u+1}, where u is the 2-adic valuation of $4k$. This answer is reminiscent of the classical image of J. We also explore some of the Toda bracket structure of the η-inverted $\mathbb{R}$-motivic stable homotopy groups.

1. Introduction

The first exotic property of motivic stable homotopy groups is that the Hopf map η is not nilpotent. This means that inverting η can be useful for understanding the global structure of motivic stable homotopy groups.

In [AM] and [GI], the η-inverted $\mathbb{C}$-motivic 2-completed stable homotopy groups $\hat{\pi}^{\mathbb{C}}_{*,*}[\eta^{-1}]$ were explicitly computed to be

$$\mathbb{F}_2[\eta^{\pm 1}][\mu, \varepsilon]/\varepsilon^2.$$

This result naturally suggests that one should study the structure of η-inverted motivic stable homotopy groups over other fields.

In the present article, we consider the η-inverted $\mathbb{R}$-motivic 2-completed stable homotopy groups $\hat{\pi}^{\mathbb{R}}_{*,*}[\eta^{-1}]$. Our main tool is the motivic Adams spectral sequence, which takes the form

$$\mathrm{Ext}_{\mathcal{A}^{\mathbb{R}}}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})[h_1^{-1}] \Rightarrow \hat{\pi}^{\mathbb{R}}_{*,*}[\eta^{-1}].$$

Here $\mathcal{A}^{\mathbb{R}}$ is the $\mathbb{R}$-motivic Steenrod algebra, and $\mathbb{M}_2^{\mathbb{R}}$ is the motivic $\mathbb{F}_2$-cohomology of $\mathbb{R}$. We will exhaustively compute this spectral sequence.

We start by computing the Adams E_2-page $\mathrm{Ext}_{\mathcal{A}^{\mathbb{R}}}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})[h_1^{-1}]$ using the ρ-Bockstein spectral sequence [H] [DI]. This spectral sequence takes the form

$$\mathrm{Ext}_{\mathcal{A}^{\mathbb{C}}}(\mathbb{M}_2^{\mathbb{C}}, \mathbb{M}_2^{\mathbb{C}})[\rho][h_1^{-1}] \Rightarrow \mathrm{Ext}_{\mathcal{A}^{\mathbb{R}}}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})[h_1^{-1}],$$

where $\mathcal{A}^{\mathbb{C}}$ is the $\mathbb{C}$-motivic Steenrod algebra and $\mathbb{M}_2^{\mathbb{C}}$ is the motivic $\mathbb{F}_2$-cohomology of $\mathbb{C}$.

The input to the ρ-Bockstein spectral sequence is completely known from [GI]. In order to deduce differentials, one first observes, as in [DI], that the groups

Date: October 30, 2015.

2000 *Mathematics Subject Classification.* 14F42, 55T15, 55Q45.

Key words and phrases. motivic homotopy theory, stable homotopy group, η-inverted stable homotopy group, Adams spectral sequence.

The first author was supported by Simons Collaboration Grant 282316. The second author was supported by NSF grant DMS-1202213.

$\mathrm{Ext}_{\mathcal{A}^{\mathbb{R}}}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})[\rho^{-1}, h_1^{-1}]$ with ρ and h_1 both inverted are easy to describe. Then there is only one pattern of ρ-Bockstein differentials that is consistent with this ρ-inverted calculation.

Having obtained the Adams E_2-page $\mathrm{Ext}_{\mathcal{A}^{\mathbb{R}}}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})[h_1^{-1}]$, the next step is to compute Adams differentials. The extension of scalars functor from $\mathbb{R}$-motivic homotopy theory to $\mathbb{C}$-motivic homotopy theory induces a map

$$\begin{array}{ccc} \mathrm{Ext}_{\mathcal{A}^{\mathbb{R}}}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})[h_1^{-1}] & \Longrightarrow & \hat{\pi}^{\mathbb{R}}_{*,*}[\eta^{-1}] \\ \downarrow & & \downarrow \\ \mathrm{Ext}_{\mathcal{A}^{\mathbb{C}}}(\mathbb{M}_2^{\mathbb{C}}, \mathbb{M}_2^{\mathbb{C}})[h_1^{-1}] & \Longrightarrow & \hat{\pi}^{\mathbb{C}}_{*,*}[\eta^{-1}] \end{array}$$

of Adams spectral sequences. The bottom Adams spectral sequence is completely understood [AM] [GI]. The Adams d_2 differentials in the top spectral sequence can then be deduced by the comparison map.

This leads to a complete description of the h_1-inverted $\mathbb{R}$-motivic Adams E_3-page. Over $\mathbb{C}$, it turns out that the h_1-inverted Adams spectral sequence collapses at this point. However, over $\mathbb{R}$, there are higher differentials that we deduce from manipulations with Massey products and Toda brackets.

In the end, we obtain an explicit description of the h_1-inverted $\mathbb{R}$-motivic Adams E_∞-page, from which we can read off the η-inverted stable motivic homotopy groups over $\mathbb{R}$.

In order to state the result, we need a bit of terminology. Because η belongs to $\hat{\pi}^{\mathbb{R}}_{1,1}$, it makes sense to use a grading that is invariant under multiplication by η. The Milnor-Witt n-stem is the direct sum $\bigoplus_p \pi^{\mathbb{R}}_{p+n,p}$. Then multiplication by η is an endomorphism of the Milnor-Witt n-stem.

Theorem 1.1.

(1) *The η-inverted Milnor-Witt 0-stem is $\mathbb{Z}_2[\eta^{\pm 1}]$, where $\mathbb{Z}_2$ is the ring of 2-adic integers.*
(2) *If $k > 1$, then the η-inverted Milnor-Witt $(4k-1)$-stem is isomorphic to $\mathbb{Z}/2^{u+1}[\eta^{\pm 1}]$ as a module over $\mathbb{Z}_2[\eta^{\pm 1}]$, where u is the 2-adic valuation of $4k$.*
(3) *The η-inverted Milnor-Witt n-stem is zero otherwise.*

For degree reasons, the product structure on $\hat{\pi}^{\mathbb{R}}_{*,*}[\eta^{-1}]$ is very simple. However, there are many interesting Toda brackets. We explore much of the 3-fold Toda bracket structure in this article. In particular, we will show that all of $\hat{\pi}^{\mathbb{R}}_{*,*}[\eta^{-1}]$ can be constructed inductively via Toda brackets, starting from just 2 and the generator of the Milnor-Witt 3-stem.

Theorem 1.1 gives a familiar answer. These groups have the same order as the classical image of J. One might expect a geometric proof that directly compares the image of J spectrum with the η-inverted $\mathbb{R}$-motivic sphere. However, the higher structure of the image of J and of the η-inverted $\mathbb{R}$-motivic sphere do not coincide. This suggests that a direct geometric proof is not possible. More specifically, let ρ_{15} be a generator of the classical image of J in the 15-stem, i.e., a class detected by $h_0^3 h_4$ in the Adams spectral sequence. Then the Toda bracket $\langle \rho_{15}, \rho_{15}, 32 \rangle$ contains a generator of the image of J in the 31-stem, i.e., is detected by $h_0^{10} h_5$

in the Adams spectral sequence. On the other hand, the analogous η-inverted $\mathbb{R}$-motivic Toda bracket does not contain a generator of the Milnor-Witt 31-stem, as shown in Remark 7.2.

The calculation of the η-inverted $\mathbb{R}$-motivic homotopy groups leads to questions about η-inverted motivic homotopy groups over other fields. We leave it to the reader to speculate on the behavior of these η-inverted groups over other fields.

2. Preliminaries

2.1. **Notation.** We continue with notation from [DI] as follows:

(1) $\mathbb{M}_2^{\mathbb{R}} = \mathbb{F}_2[\tau, \rho]$ is the motivic cohomology of $\mathbb{R}$ with $\mathbb{F}_2$ coefficients, where τ and ρ have bidegrees $(0,1)$ and $(1,1)$, respectively.
(2) $\mathbb{M}_2^{\mathbb{C}} = \mathbb{F}_2[\tau]$ is the motivic cohomology of $\mathbb{C}$ with $\mathbb{F}_2$ coefficients, where τ has bidegree $(0,1)$.
(3) $\mathcal{A}^{\mathbb{R}}$ is the mod 2 motivic Steenrod algebra over $\mathbb{R}$.
(4) $\mathcal{A}^{\mathbb{C}}$ is the mod 2 motivic Steenrod algebra over $\mathbb{C}$.
(5) $\mathcal{A}^{\mathrm{cl}}$ is the classical mod 2 Steenrod algebra.
(6) $\mathrm{Ext}_{\mathbb{R}}$ is the trigraded ring $\mathrm{Ext}_{\mathcal{A}^{\mathbb{R}}}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})$.
(7) $\mathrm{Ext}_{\mathbb{C}}$ is the trigraded ring $\mathrm{Ext}_{\mathcal{A}^{\mathbb{C}}}(\mathbb{M}_2^{\mathbb{C}}, \mathbb{M}_2^{\mathbb{C}})$.
(8) $\mathrm{Ext}_{\mathrm{cl}}$ is the trigraded ring $\mathrm{Ext}_{\mathcal{A}^{\mathrm{cl}}}(\mathbb{F}_2, \mathbb{F}_2)$.
(9) $\mathcal{R} = \mathbb{F}_2[\rho, h_1^{\pm 1}]$.
(10) $\hat{\pi}^{\mathbb{C}}_{*,*}$ is the motivic stable homotopy ring of the 2-completed motivic sphere spectrum over $\mathbb{C}$.
(11) $\hat{\pi}^{\mathbb{R}}_{*,*}$ is the motivic stable homotopy ring of the 2-completed motivic sphere spectrum over $\mathbb{R}$.
(12) The symbols v_1^4 and P are used interchangeably for the Adams periodicity operator.

2.2. **Grading conventions.** We follow [I] in grading Ext according to (s, f, w), where:

(1) f is the Adams filtration, i.e., the homological degree.
(2) $s + f$ is the internal degree, i.e., corresponds to the first coordinate in the bidegree of the Steenrod algebra.
(3) s is the stem, i.e., the internal degree minus the Adams filtration.
(4) w is the weight.

We will consider the groups $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$ in which h_1 has been inverted. The degree of h_1 is $(1,1,1)$. As in [GI], for this purpose it is convenient to introduce the following gradings whose values are zero for h_1.

(5) $mw = s - w$ is the Milnor-Witt degree.
(6) $c = s + f - 2w$ is the Chow degree.

In order to avoid notational clutter, we will drop h_1 from the notation, except in the formal statements of the main results. Since h_1 is a unit, no information is lost by doing this. The correct powers of h_1 can always be recovered by checking degrees.

For example, in Lemma 3.1 below, we claim that there is a differential $d_3^{\rho}(v_1^4) = \rho^3 v_2$ in the ρ-Bockstein spectral sequence. Strictly speaking, this formula is nonsensical because $d_3^{\rho}(v_1^4)$ has Adams filtration 5 while v_2 has Adams filtration 1. The correct full formula is $d_3^{\rho}(v_1^4) = \rho^3 h_1^4 v_2$.

If we are to ignore multiples of h_1, we must rely on gradings that take value 0 on h_1. This explains our preference for Milnor-Witt degree mw and Chow degree c.

3. The ρ-Bockstein spectral sequence

Recall [H, DI] that the ρ-Bockstein spectral sequence takes the form

$$\mathrm{Ext}_{\mathbb{C}}[\rho] \Rightarrow \mathrm{Ext}_{\mathbb{R}}\,.$$

After inverting h_1, by [GI, Theorem 1.1] this takes the form

$$\mathcal{R}[v_1^4, v_2, v_3, \dots] \Rightarrow \mathrm{Ext}_{\mathbb{R}}[h_1^{-1}],$$

where $\mathcal{R} = \mathbb{F}_2[\rho, h_1^{\pm 1}]$. Table 1 lists the generators of the Bockstein E_1-page.

Table 1. Bockstein E_1-page generators

(mw, c)	generator
$(0,1)$	ρ
$(4,4)$	v_1^4
$(3,1)$	v_2
$(7,1)$	v_3
$(15,1)$	v_4
$(2^n-1,1)$	v_n

Lemma 3.1. *In the ρ-Bockstein spectral sequence, there are differentials $d^{\rho}_{2^n-1}(v_1^{2^n}) = \rho^{2^n-1}v_n$ for $n \geq 2$. All other non-zero differentials follow from the Leibniz rule.*

The first few examples of these differentials are $d_3(v_1^4) = \rho^3 v_2$, $d_7(v_1^8) = \rho^7 v_3$, and $d_{15}(v_1^{16}) = \rho^{15} v_4$.

Proof. Inverting ρ induces a map

$$\begin{array}{ccc} \mathrm{Ext}_{\mathbb{C}}[h_1^{-1}][\rho] & \xRightarrow{\rho\text{-Bss}} & \mathrm{Ext}_{\mathbb{R}}[h_1^{-1}] \\ \downarrow{\scriptstyle \rho\text{-inv}} & & \downarrow{\scriptstyle \rho\text{-inv}} \\ \mathrm{Ext}_{\mathbb{C}}[h_1^{-1}][\rho^{\pm 1}] & \xRightarrow{\rho\text{-Bss}} & \mathrm{Ext}_{\mathbb{R}}[h_1^{-1}, \rho^{-1}] \end{array}$$

of ρ-Bockstein spectral sequences. We will establish differentials in the ρ-inverted spectral sequence. The map of spectral sequences then implies that the same differentials occur when ρ is not inverted.

Recall [DI, Theorem 4.1] that there is an isomorphism $\mathrm{Ext}_{\mathrm{cl}}[\rho^{\pm 1}] \cong \mathrm{Ext}_{\mathbb{R}}[\rho^{-1}]$ sending the classical element h_0 to the motivic element h_1. Using also that $\mathrm{Ext}_{\mathrm{cl}}[h_0^{-1}] = \mathbb{F}_2[h_0^{\pm 1}]$, it follows that $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}, \rho^{-1}]$ is isomorphic to $\mathcal{R}[\rho^{-1}]$. Then the ρ-inverted ρ-Bockstein spectral sequence takes the form

$$\mathcal{R}[\rho^{-1}][v_1^4, v_2, v_3, \dots] \xRightarrow{\rho\text{-Bss}} \mathcal{R}[\rho^{-1}].$$

Because the target of the ρ-inverted spectral sequence is very small, essentially everything must either support a differential or be hit by a differential.

The ρ-Bockstein differentials have degree $(-1,0)$ with respect to the grading (mw,c) used in Table 1. The elements $\rho^k v_2$ cannot support differentials because there are no elements in the Milnor-Witt 2-stem. The only possibility is that after inverting ρ, there is a ρ-Bockstein differential $d_3(v_1^4)=\rho^3 v_2$.

Then the ρ-inverted E_4-page is $\mathcal{R}[v_1^8, v_3, v_4, \ldots]$. The elements $\rho^k v_3$ cannot support differentials because the ρ-inverted E_4-page has no elements in Milnor-Witt 6-stem. The only possibility is that after inverting ρ, there is a ρ-Bockstein differential $d_7(v_1^8)=\rho^7 v_3$.

In general, the ρ-inverted $E_{2^{n-1}}$-page is $\mathcal{R}[v_1^{2^n}, v_n, v_{n+1}, \ldots]$. The elements $\rho^k v_n$ cannot support differentials because the ρ-inverted $E_{2^{n-1}}$-page has no elements in the Milnor-Witt (2^n-2)-stem. The only possibility is that after inverting ρ, there is a ρ-Bockstein differential $d_{2^n-1}(v_1^{2^n})=\rho^{2^n-1}v_n$. ∎

The ρ-Bockstein E_∞-page can be directly computed from the differentials in Lemma 3.1. We write P rather than v_1^4.

Proposition 3.2. *The ρ-Bockstein E_∞-page is the $\mathcal{R}$-algebra on the generators $P^{2^{n-1}k}v_n$ for $n\geq 2$ and $k\geq 0$ (see Table 2), subject to the relations*

$$\rho^{2^n-1}P^{2^{n-1}k}v_n = 0$$

for $n\geq 2$ and $k\geq 0$, and

$$P^{2^{n-1}k}v_n \cdot P^{2^{m-1}j}v_m + P^{2^{n-1}(k+2^{m-n}j)}v_n \cdot v_m = 0$$

for $m\geq n\geq 2$, $k\geq 0$, and $j\geq 0$.

TABLE 2. Bockstein E_∞-page generators

(mw,c)	generator	ρ-torsion
$(0,1)$	ρ	∞
$(0,0)$	h_1	∞
$(3,1)+k(8,8)$	$P^{2k}v_2$	3
$(7,1)+k(16,16)$	$P^{4k}v_3$	7
$(15,1)+k(32,32)$	$P^{8k}v_4$	15
$(2^n-1,1)+k(2^{n+1},2^{n+1})$	$P^{2^{n-1}k}v_n$	2^n-1

Remark 3.3. In practice, the relations mean that all P's can be shifted onto the v_n with minimal n in any monomial. Thus an $\mathcal{R}$-module basis is given by monomials of the form $P^{2^{n-1}k}v_n\cdot v_{m_1}\cdots v_{m_a}$, where $n\leq m_1\leq\cdots\leq m_a$. For example,

$$P^2v_2\cdot P^4v_2 = P^6v_2\cdot v_2$$
$$P^4v_2\cdot P^8v_3 = P^{12}v_2\cdot v_3$$
$$P^4v_3\cdot P^{48}v_5 = P^{52}v_3\cdot v_5.$$

4. The Adams E_2-page

Having obtained the ρ-Bockstein E_∞-page in Section 3, our next task is to consider hidden extensions in $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$. We will show that what you see (in the Bockstein E_∞ page) is what you get (in Ext), as there are no hidden relations. This will require some careful analysis of degrees, as well as some manipulations with Massey products.

Lemma 4.1. *For each $n \geq 2$ and $k \geq 0$, the element $P^{2^{n-1}k}v_n$ of the Bockstein E_∞-page detects a unique element of* $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$.

Proof. We need to show that $P^{2^{n-1}k}v_n$ does not share bidegree with an element that is divisible by ρ.

First suppose that $P^{2^{n-1}k}v_n$ has the same bidegree as $\rho^b P^{2^{m-1}j}v_m$. Then

$$(2^n - 1, 1) + k(2^{n+1}, 2^{n+1}) = (2^m - 1, 1) + j(2^{m+1}, 2^{m+1}) + b(0, 1).$$

Considering only the Milnor-Witt degree, we have

$$2^n(2k+1) = 2^m(2j+1).$$

Therefore, $n = m$ and $k = j$, so $b = 0$.

Now suppose that $P^{2^{n-1}k}v_n$ shares bidegree with some element x. By Remark 3.3, we may assume that x is of the form $\rho^b P^{2^{m_1-1}j}v_{m_1} \cdot v_{m_2} \cdots v_{m_a}$, where $m_1 \leq m_2 \leq \cdots \leq m_a$. We may also assume that $b \leq 2^{m_1} - 2$, since $\rho^{2^{m_1}-1}P^{2^{m-1}j}v_{m_1} = 0$. Because of the previous paragraph, we may assume that $a \geq 2$. We wish to show that $b = 0$.

We first show that $n \geq m_a$. Let $u(x)$ be the difference $mw - c$. We have $u(P^{2^{m_1-1}j}v_{m_1}) = 2^{m_1} - 2$ and $u(\rho) = -1$. Since $b \leq 2^{m_1} - 2$, it follows that $u(\rho^b P^{2^{m_1-1}j}v_{m_1}) \geq 0$. Thus

$$2^n - 2 = u(P^{2^{n-1}k}v_n) = u(\rho^b P^{2^{m_1-1}j}v_{m_1}) + u(v_{m_2} \cdots v_{m_a}) \geq u(v_{m_a}) = 2^{m_a} - 2,$$

so that $n \geq m_a$.

Now consider the Milnor-Witt and Chow degrees modulo 4. We have

$$(-1, 1) \equiv (-a, a + b) \pmod 4,$$

so $a \equiv 1 \pmod 4$ and $b \equiv 0 \pmod 4$. Thus either $b = 0$, which was what we wanted to show, or $b \geq 4$.

We may now assume that $b \geq 4$. Since $\rho^4 P^{2j}v_2 = 0$, we must have $m_1 \geq 3$, so that all m_i, and also n, are at least 3.

Next, consider degrees modulo 8. Comparing degrees gives

$$(-1, 1) \equiv (-a, a + b) \pmod 8.$$

Thus $b \equiv 0 \pmod 8$, so that $b \geq 8$. Since $\rho^8 P^{4j}v_3 = 0$, we must have that j_1, and therefore also n and all other j_i, to be at least 4. This argument can be continued to establish that b and n must be arbitrarily large under the assumption that $b > 0$. ∎

Remark 4.2. Lemma 4.1 allows us to use unambiguously the same notation $P^{2^{n-1}k}v_n$ for an element of $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$.

Lemma 4.3. *For each $n \geq 2$ and $k \geq 0$, the element $P^{2^{n-1}k}v_n \cdot v_n$ of the Bockstein E_∞-page detects a unique element of* $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$.

Proof. The Milnor-Witt degree of $P^{2^{n-1}k}v_n \cdot v_n$ is even, while the Milnor-Witt degree of $\rho^b P^{2^{m-1}j}v_m$ is odd. Therefore, these elements cannot share bidegree.

Now suppose that the element $P^{2^{n-1}k}v_n \cdot v_n$ has the same bidegree as the element $\rho^b P^{2^{m_1-1}j}v_{m_1} \cdot v_{m_2} \cdots v_{m_a}$, with $m_1 \leq m_2 \leq \cdots \leq m_a$, $b \leq 2^{m_1} - 2$, and $a \geq 2$. The rest of the proof is essentially the same as the proof of Lemma 4.1. Consider $u = mw - c$ to get that $n \geq m_a$. Then consider congruences $(-2, 2) \equiv (-a, a + b)$ modulo higher and higher powers of 2 to obtain that $b = 0$. ∎

Remark 4.4. The obvious generalization of Lemma 4.3 to elements of the form $P^{2^{n-1}k}v_n \cdot v_m$ is false. For example, $P^2v_2 \cdot v_5$ has the same degree as $\rho^4 v_3^6$.

Remark 4.5. Lemmas 4.1 and 4.3 are equivalent to the claim that there are no ρ multiples in the ρ-Bockstein E_∞-page in the same bidegrees as $P^{2^{n-1}k}v_n$ or $P^{2^{n-1}k}v_n \cdot v_n$. This implies that there are also no ρ multiples in $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$ that share bidegree with these elements; we will need this fact later. The point is that the ρ-Bockstein spectral sequence does not allow for hidden extensions by ρ.

Lemma 4.6. $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$ *is zero when the Milnor-Witt stem mw and the Chow degree c are both equal to* $2i$ *with* $i \geq 1$.

Proof. Under the condition $mw = c = 2i$, inspection of Table 1 shows that the ρ-Bockstein E_1-page consists of products of elements of the form v_1^4 or $\rho^{2^n+2^m-4}v_nv_m$. In the E_∞-page, $\rho^{2^n+2^m-4}v_nv_m = 0$ since $\rho^{2^n-1}v_n = 0$. Also, v_1^{4k} supports a differential for all $k \geq 0$. ∎

Lemma 4.7. *For each* $n \geq 2$, $k \geq 1$ *and* $m > n$, *we have a Massey product*

$$P^{2^{n-1}k+2^{m-2}}v_n = \langle \rho^{2^m-2^n}v_m, \rho^{2^n-1}, P^{2^{n-1}k}v_n \rangle$$

in $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$ *with no indeterminacy.*

Proof. The Massey product formula follows from the Bockstein differential $d_{2^m-1}^{\rho}(P^{2^{m-2}}) = \rho^{2^m-1}v_m$ and May's convergence theorem [M1, Theorem 4.1]. There are no crossing Bockstein differentials as all classes are in nonnegative ρ-filtration.

The indeterminacy of this bracket is generated by products of the form $\rho^{2^m-2^n}v_m \cdot x$ and $y \cdot P^{2^{n-1}k}v_n$, where x and y have appropriate bidegrees. We showed in Lemma 4.6 that 0 is the only possibility for x or y. ∎

Remark 4.8. Lemma 4.7 gives many different Massey products for the same element. For example,

$$P^8v_2 = \langle \rho^4 v_3, \rho^3, P^6v_2 \rangle$$
$$P^8v_2 = \langle \rho^{12} v_4, \rho^3, P^4v_2 \rangle$$
$$P^8v_2 = \langle \rho^{28} v_5, \rho^3, v_2 \rangle.$$

May's convergence theorem only says that these Massey products are detected by the same element of the ρ-Bockstein E_∞-page. By Lemma 4.1, this E_∞-page element detects just one element of $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$, so all of the Massey products must in fact be equal.

Lemma 4.9. *For* $m > n \geq 2$, *there is a Massey product*

$$P^{2^{n-1}k+2^{m-2}}v_n = \langle P^{2^{n-1}k}v_n, \rho^{2^m-2}v_m, \rho \rangle$$

in $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$ *with no indeterminacy.*

Proof. The Massey product formula follows from the Bockstein differential $d_{2^m-1}^{\rho}(P^{2^{m-2}}) = \rho^{2^m-1}v_m$ and May's convergence theorem [M1, Theorem 4.1]. There are no crossing Bockstein differentials as all classes are in nonnegative ρ-filtration.

The indeterminacy of this bracket is generated by products of the form $P^{2^{n-1}k}v_n \cdot x$ and $y \cdot \rho$. We showed in Lemma 4.6 that 0 is the only possibility for x. We observed

in Remark 4.5 that $y \cdot \rho$ must be zero because there are no multiples of ρ in the appropriate bidegree. ∎

The relations in the Bockstein E_∞-page given in Proposition 3.2 may lift to $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$ with additional terms that are multiples of ρ. In other words, there may be hidden relations in the Bockstein spectral sequence. For example, for degree reasons it is possible that $P^2 v_2 \cdot P^{16} v_5 + P^{18} v_2 \cdot v_5$ equals $\rho^4 P^{16} v_3 \cdot v_3^5$. Proposition 4.10 shows that there are no such hidden terms in the relations in $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$.

Proposition 4.10. *There are no hidden relations in the Bockstein spectral sequence.*

Proof. There can be no hidden ρ extensions in the ρ-Bockstein spectral sequence, so we need only compute the products $P^{2^{n-1}k} v_n \cdot P^{2^{m-1}j} v_m$ in $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$ for $m \geq n$.

Lemma 4.7 implies that $P^{2^{n-1}k} v_n \cdot P^{2^{m-1}j} v_m$ equals

$$P^{2^{n-1}k} v_n \langle \rho^{2^m} v_{m+1}, \rho^{2^m - 1}, P^{2^{m-1}(j-1)} v_m \rangle.$$

Shuffle to obtain

$$\langle P^{2^{n-1}k} v_n, \rho^{2^m} v_{m+1}, \rho^{2^m - 1} \rangle P^{2^{m-1}(j-1)} v_m.$$

This expression is contained in

$$\langle P^{2^{n-1}k} v_n, \rho^{2^{m+1} - 2} v_{m+1}, \rho \rangle P^{2^{m-1}(j-1)} v_m,$$

which equals $P^{2^{n-1}k + 2^{m-1}} v_n \cdot P^{2^{m-1}(j-1)} v_m$ by Lemma 4.9.

By induction, $P^{2^{n-1}k} v_n \cdot P^{2^{m-1}j} v_m$ equals $P^{2^{n-1}(k + 2^{m-n}j)} v_n \cdot v_m$. ∎

Theorem 4.11. $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$ *is the $\mathcal{R}$-algebra on the generators $P^{2^{n-1}k} v_n$ for $n \geq 2$ and $k \geq 0$ (see Table 2), subject to the relations*

$$\rho^{2^n - 1} P^{2^{n-1}k} v_n = 0$$

for $n \geq 2$ and $k \geq 0$, and

$$P^{2^{n-1}k} v_n \cdot P^{2^{m-1}j} v_m + P^{2^{n-1}(k + 2^{m-n}j)} v_n \cdot v_m = 0$$

for $m \geq n \geq 2$, $k \geq 0$, and $j \geq 0$.

Proof. This follows immediately from Propositions 3.2 and 4.10. ∎

Remark 4.12. Analogously to Remark 3.3, an $\mathcal{R}$-module basis for $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$ is given by monomials of the form $P^{2^{n-1}k} v_n \cdot v_{m_1} \cdots v_{m_a}$, where $n \leq m_1 \leq \cdots \leq m_a$.

5. Adams differentials

Before computing with the h_1-inverted $\mathbb{R}$-motivic Adams spectral sequence, we will consider convergence. A priori, there could be an infinite family of homotopy classes linked together by infinitely many hidden η multiplications. These classes would not be detected in $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$. Lemma 5.1 implies that this cannot occur for degree reasons.

Lemma 5.1. *Let $m > 0$ be a fixed Milnor-Witt stem. There exists a constant A such that $\mathrm{Ext}_{\mathbb{R}}^{(s,f,w)}$ vanishes when $s - w = m$, $s \neq 0$, $f > A$, and $f > s + 1$.*

Lemma 5.1 can be restated in the following more casual form. Within a fixed Milnor-Witt stem, there exists a horizontal line and a line of slope 1 such that $\mathrm{Ext}_{\mathbb{R}}$ vanishes in the region above both lines, except in the 0-stem. Figure 1 depicts the shape of the vanishing region.

FIGURE 1. The vanishing region in a Milnor-Witt stem



Proof. This argument occurs in $\mathrm{Ext}_{\mathbb{R}}$, where h_1 has not been inverted.

As explained in [DI, Theorem 4.1], the elements in the m-stem of the classical Ext groups $\mathrm{Ext}_{\mathrm{cl}}$ correspond to elements of $\mathrm{Ext}_{\mathbb{R}}$ in the Milnor-Witt m-stem that remain non-zero after ρ is inverted, i.e., that support infinitely many multiplications by ρ. Each stem of $\mathrm{Ext}_{\mathrm{cl}}$ is finite except for the 0-stem. For $m > 0$, choose A to be larger than the Adams filtrations of all of the elements in the m-stem of $\mathrm{Ext}_{\mathrm{cl}}$. Then A is larger than the Adams filtrations of every element of $\mathrm{Ext}_{\mathbb{R}}$ in the Milnor-Witt m-stem that remain non-zero after ρ is inverted.

Let x be a non-zero element of $\mathrm{Ext}_{\mathbb{R}}^{(s,f,w)}$ such that $s - w = m$, $f > A$, and $f > s + 1$. We will show that s must equal zero.

The choice of A guarantees that x is annihilated by some positive power of ρ. Suppose that $\rho^k x = 0$ but $\rho^{k-1}x$ is non-zero, for some $k > 0$. Then there must be a differential in the ρ-Bockstein spectral sequence of the form $d_k(y) = \rho^k x$, where y is an element of $\mathrm{Ext}_{\mathbb{C}}$ in degree $(s - k + 1, f - 1, w - k)$.

The argument from [A] establishes a vanishing line of slope 1 in $\mathrm{Ext}_{\mathbb{C}}$. The conditions $s > f + 1$ and $k > 0$ imply that the element y lies strictly above this vanishing line, so it must be of the form $\tau^a h_0^b$ with $b \geq 1$. The only ρ-Bockstein differentials on such classes are $d_1(\tau^{2c+1}h_0^b) = \rho\tau^{2c}h_0^{b+1}$, which implies that x must be of the form $\tau^{2c}h_0^b$. This shows that $s = 0$. ∎

The h_1-inverted motivic Adams spectral sequence over $\mathbb{C}$ was studied in [GI, AM]. It takes the form

$$\mathbb{F}_2[h_1^{\pm 1}, P, v_2, v_3, \dots] \Rightarrow \hat{\pi}_{*,*}^{\mathbb{C}}[\eta^{-1}],$$

where $\hat{\pi}_{*,*}^{\mathbb{C}}[\eta^{-1}]$ is the η-inverted motivic stable homotopy ring of the 2-completed motivic sphere spectrum over $\mathbb{C}$. This spectral sequence has differentials

$$d_2(P^k v_n) = P^k v_{n-1}^2$$

for all $k \geq 0$ and all $n \geq 3$.

Lemma 5.2. *In the h_1-inverted $\mathbb{R}$-motivic Adams spectral sequence, there are differentials*

$$d_2(P^{2^{n-1}k}v_n) = P^{2^{n-1}k}v_{n-1}^2$$

for all $k \geq 0$ and all $n \geq 3$.

Proof. There is an extension of scalars functor from $\mathbb{R}$-motivic homotopy theory to $\mathbb{C}$-motivic homotopy theory. This functor induces a map

$$\begin{array}{ccc} \mathrm{Ext}_{\mathcal{A}^{\mathbb{R}}}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})[h_1^{-1}] & \Longrightarrow & \hat{\pi}_{*,*}^{\mathbb{R}}[\eta^{-1}] \\ \downarrow & & \downarrow \\ \mathrm{Ext}_{\mathcal{A}^{\mathbb{C}}}(\mathbb{M}_2^{\mathbb{C}}, \mathbb{M}_2^{\mathbb{C}})[h_1^{-1}] & \Longrightarrow & \hat{\pi}_{*,*}^{\mathbb{C}}[\eta^{-1}] \end{array}$$

from the $\mathbb{R}$-motivic Adams spectral sequence to the $\mathbb{C}$-motivic Adams spectral sequence. This map takes ρ to zero.

The map of spectral sequences implies that the $\mathbb{R}$-motivic Adams differential $d_2(P^{2^{n-1}k}v_n)$ equals $P^{2^{n-1}k}v_{n-1}^2$ plus terms that are divisible by ρ. Lemma 4.3 implies that there are no possible additional terms in the relevant bidegree. ∎

We record the following simple computation, as we will employ it several times.

Lemma 5.3. *Let S be an $\mathbb{F}_2$-algebra. Let $B = S[w_1, w_2, \dots]$ be a polynomial ring in infinitely many variables, and define a differential on B by $\partial(w_n) = w_{n-1}^2$ for $n \geq 2$. Then $\mathrm{H}^*(B, \partial) \cong S[w_1]/w_1^2$.*

Lemma 5.3 implies, for example, that the h_1-inverted $\mathbb{C}$-motivic Adams E_3-page is $\mathbb{F}_2[h_1^{\pm 1}, P, v_2]/v_2^2$.

Proposition 5.4. *The h_1-inverted $\mathbb{R}$-motivic Adams E_3-page is free as an $\mathcal{R}$-module on the generators listed in Table 3 for $n \geq 2$, $k \geq 0$, and $j \geq 0$. Almost all products of these generators are zero, except that*

$$P^{4k}v_2 \cdot P^{4j+2}v_2 = P^{4k+4j+2}v_2^2$$

and

$$\rho^{2^{n-1}-1}P^{2^{n-1}\cdot 2k}v_n \cdot \rho^{2^{n-1}-1}P^{2^{n-1}(2j+1)}v_n = \rho^{2^n-2}P^{2^{n-1}(2k+2j+1)}v_n^2$$

for $n \geq 3$.

TABLE 3. $\mathcal{R}$-module generators for the Adams E_3-page

(mw, c)	generator	ρ-torsion
$(0,0)$	1	∞
$(3,1) + k(8,8)$	$P^{2k}v_2$	3
$(7,4) + k(16,16)$	$\rho^3 P^{4k}v_3$	4
$(15,8) + k(32,32)$	$\rho^7 P^{8k}v_4$	8
$(2^n - 1, 2^{n-1}) + k(2^{n+1}, 2^{n+1})$	$\rho^{2^{n-1}-1}P^{2^{n-1}k}v_n$	2^{n-1}
$(6,2) + (2j+1)(8,8)$	$P^{2(2j+1)}v_2^2$	3
$(14,2) + (2j+1)(16,16)$	$P^{4(2j+1)}v_3^2$	7
$(30,2) + (2j+1)(32,32)$	$P^{8(2j+1)}v_4^2$	15
$(2^{n+1} - 2, 2) + (2j+1)(2^{n+1}, 2^{n+1})$	$P^{2^{n-1}(2j+1)}v_n^2$	$2^n - 1$

Remark 5.5. The relations in Proposition 5.4 are just the ones that are obvious from the notation. For example,

$$v_2 \cdot P^2 v_2 = P^2 v_2^2$$
$$\rho^3 P^4 v_3 \cdot \rho^3 P^8 v_3 = \rho^6 P^{12} v_3^2.$$

Proof. Let $\mathrm{Ext}\langle k, b\rangle$ be the $\mathbb{F}_2[h_1^{\pm 1}]$-submodule of the h_1-inverted $\mathbb{R}$-motivic Adams E_2-page on generators of the form $\rho^b P^k v_{m_1} v_{m_2} \cdots v_{m_a}$ such that $m_1 \leq m_2 \leq \cdots \leq m_a$. Note that $b \leq 2^{m_1} - 2$ in this situation, since $\rho^{2^{m_1}-1} P^k v_{m_1} = 0$. Also, k must be a multiple of 2^{m_1-1}. By Lemma 5.2, each $\mathrm{Ext}\langle k, b\rangle$ is a differential graded submodule. Thus it suffices to compute the cohomology of each $\mathrm{Ext}\langle k, b\rangle$.

We start with $\mathrm{Ext}\langle 0, b\rangle$, which is equal to to $\rho^b \cdot \mathbb{F}_2[h_1^{\pm 1}, v_m, v_{m+1}, \ldots]$ as a differential graded $\mathbb{F}_2[h_1^{\pm 1}]$-module, where m is the smallest integer such that $b \leq 2^m - 2$. Now Lemma 5.3 implies that $H^*(\mathrm{Ext}\langle 0, b\rangle, d_2)$ is a free $\mathbb{F}_2[h_1^{\pm 1}]$-module on two generators ρ^b and $\rho^b v_m$.

So far, we have demonstrated that the powers of ρ and the elements

$$v_2, \rho v_2, \rho^2 v_2, \rho^3 v_3, \ldots \rho^6 v_3, \rho^7 v_4, \ldots$$

are present in the h_1-inverted $\mathbb{R}$-motivic Adams E_3-page.

The module $\mathrm{Ext}\langle k, b\rangle$ is zero when k is odd.

Now assume that k is equal to 2 modulo 4. If $b \leq 2$, then $\mathrm{Ext}\langle k, b\rangle$ is equal to $\rho^b P^k v_2 \cdot \mathbb{F}_2[h_1^{\pm 1}, v_2, v_3, \ldots]$ as a differential graded $\mathbb{F}_2[h_1^{\pm 1}]$-module. Lemma 5.3 implies that $H^*(\mathrm{Ext}\langle k, b\rangle, d_2)$ is a free $\mathbb{F}_2[h_1^{\pm 1}]$-module on two generators $\rho^b P^k v_2$ and $\rho^b P^k v_2^2$. If $b \geq 3$, then $\mathrm{Ext}\langle k, b\rangle$ is zero because $\rho^3 P^k v_2 = 0$.

We have now shown that the elements

$$P^k v_2, \rho P^k v_2, \rho^2 P^k v_2, P^k v_2^2, \rho P^k v_2^2, \rho^2 P^k v_2^2$$

are present in the h_1-inverted $\mathbb{R}$-motivic Adams E_3-page for all k congruent to 2 modulo 4.

Next assume that k is equal to 4 modulo 8. If $b \leq 2$, then $\mathrm{Ext}\langle k, b\rangle$ is the free $\mathbb{F}_2[h_1^{\pm 1}]$-module on generators $\rho^b P^k v_{m_1} \cdots v_{m_a}$ such that m_1 equals 2 or 3, and $m_1 \leq \cdots \leq m_a$. There is a short exact sequence

$$\mathrm{Ext}\langle k, b\rangle \to \rho^b P^k \cdot \mathbb{F}_2[h_1^{\pm 1}, v_2, v_3, \ldots] \to \rho^b P^k \cdot \mathbb{F}_2[h_1^{\pm 1}, v_4, v_5, \ldots].$$

By Lemma 5.3, the homology of the middle term has two generators $\rho^b P^k$ and $\rho^b P^k v_2$, while the homology of the right term has two generators $\rho^b P^k$ and $\rho^b P^k v_4$. Analysis of the long exact sequence in homology shows that $H^*(\mathrm{Ext}\langle k, b\rangle, d_2)$ has two generators $\rho^b P^k v_2$ and $\rho^b P^k v_3^2$.

Now assume that $3 \leq b \leq 6$. Since $\rho^b P^k v_2 = 0$, we get that $\mathrm{Ext}\langle k, b\rangle$ is equal to $\rho^b P^k v_3 \cdot \mathbb{F}_2[h_1^{\pm 1}, v_3, v_4, \ldots]$. Lemma 5.3 implies that $H^*(\mathrm{Ext}\langle k, b\rangle, d_2)$ is a free $\mathbb{F}_2[h_1^{\pm 1}]$-module on two generators $\rho^b P^k v_3$ and $\rho^b P^k v_3^2$.

Finally, if $b \geq 7$, then $\mathrm{Ext}\langle k, b\rangle$ is zero because $\rho^7 P^k v_2 = 0$ and $\rho^7 P^k v_3 = 0$. This finishes the argument when k is equal to 4 modulo 8, and we have shown that $\mathrm{Ext}_{\mathbb{R}}[h_1^{\pm 1}]$ contains the elements

$$P^k v_2, \rho P^k v_2, \rho^2 P^k v_2$$
$$\rho^3 P^k v_3, \ldots, \rho^6 P^k v_3$$
$$P^k v_3^2, \rho P^k v_3^2, \ldots, \rho^6 P^k v_3^2.$$

Analysis of the other cases is the same as the argument for $k = 4$ modulo 8. The details depend on the value of k modulo 2^i and inequalities of the form $2^j - 1 \leq b \leq 2^{j+1} - 2$. In each case there is a short exact sequence of differential graded modules whose first term is $\mathrm{Ext}\langle k, b\rangle$ and whose other two terms have homology that is computed by Lemma 5.3. ∎

We have now calculated the h_1-inverted $\mathbb{R}$-motivic E_3-page. This E_3-page is displayed in Figure 2. Beware that the grading on this chart is not the same as in a standard Adams chart. The Milnor-Witt stem $mw = s - w$ is plotted on the horizontal axis, while the Chow degree $c = s + f - 2w$ is plotted on the vertical axis. As a result, an Adams d_r differential has slope $-r + 1$, rather than slope $-r$. Vertical lines in Figure 2 represent multiplications by ρ.

Our next goal is to establish the Adams d_3 differentials. Inspection of Figure 2 reveals that the only possible non-zero d_3 differentials might be supported on elements of the form $\rho^b P^{2^{n-1}k} v_n$ for $n \geq 4$. In fact, these differentials all occur, as indicated in Figure 2 by lines that go left one unit and up two units. We will establish these d_3 differentials by first proving a homotopy relation in Lemma 5.6.

Lemma 5.6. *For each $n \geq 2$ and $j \geq 0$, the element $P^{2^{n-1}(2j+1)} v_n^2$ is a permanent cycle that detects a ρ-divisible element of the η-inverted $\mathbb{R}$-motivic homotopy groups.*

Proof. Inspection of Figure 2 shows that $P^{2^{n-1}(2j+1)} v_n^2$ cannot support a differential.

Lemma 4.9 implies that $P^{2^{n-1}(2j+1)} v_n^2$ belongs to the Massey product $\langle \rho, \rho^{2^{n+1}-2} v_{n+1}, P^{2^n j} v_n^2 \rangle$ in $\mathrm{Ext}_{\mathbb{R}}[h_1^{-1}]$. In fact, the Massey product has no indeterminacy because of Remark 4.5 and Lemma 4.6.

We will now apply Moss's convergence theorem [M3, Theorem 1.2] to this Massey product. There is an Adams differential $d_2(P^{2^n j} v_{n+1}) = P^{2^n j} v_n^2$, so $P^{2^n j} v_n^2$ detects the homotopy element 0. By inspection of Figure 2, $\rho^{2^{n+1}-2} v_{n+1}$ is a permanent cycle; let α be a homotopy element detected by it. Moreover, $\rho\alpha$ is zero in homotopy because there are no classes in higher filtration that could detect it.

Moss's convergence theorem says that the Toda bracket $\langle \rho, \alpha, 0\rangle$ contains an element that is detected by $P^{2^{n-1}(2j+1)} v_n^2$. This Toda bracket consists entirely of multiples of ρ. ∎

Lemma 5.7.

$$d_3(\rho^{2^{n-1}-1} P^{2^{n-1}k} v_n) = P^{2^{n-3}+2^{n-1}k} v_{n-2}^2$$

for $n \geq 4$.

Proof. Lemma 5.6 shows that $P^{2^{n-3}+2^{n-1}k} v_{n-2}^2$ detects a class that is divisible by ρ. By inspection of Figure 2, there are no classes in lower filtration. Therefore, $P^{2^{n-3}+2^{n-1}k} v_{n-2}^2$ must detect zero, i.e., must be hit by a differential. It is apparent from Figure 2 that there is only one possible differential. ∎

Lemma 5.8 describes the higher Adams differentials.

Lemma 5.8. *For $n \geq r + 1$ and $r \geq 3$,*

$$d_r(\rho^{2^n - 2^{n-r+2} - r + 2} P^{2^{n-1}k} v_n) = P^{2^{n-1}k + 2^{n-2} - 2^{n-r}} v_{n-r+1}^2.$$

Proof. The proof is essentially the same as the proof of Lemma 5.7. In Milnor-Witt stem congruent to 2 modulo 4, Lemma 5.6 implies that every homotopy element is divisible by ρ. This implies that they must all be hit by differentials. Figure 2 indicates that there is just one possible pattern of differentials. ∎

From Lemma 5.8, it is straightforward to derive the h_1-inverted Adams E_∞-page, as shown in Figure 3.

Proposition 5.9. *The h_1-inverted Adams E_∞-page is the $\mathcal{R}$-module on generators given in Table 4 for $n \geq 2$.*

TABLE 4. $\mathcal{R}$-module generators for the Adams E_∞-page

(mw, c)	generator	ρ-torsion
$(0,0)$	1	∞
$(3,1)+k(8,8)$	$P^{2k}v_2$	3
$(7,4)+k(16,16)$	$\rho^3 P^{4k}v_3$	4
$(15,11)+k(32,32)$	$\rho^{10}P^{8k}v_4$	5
$(2^n-1, 2^n-n-1)+k(2^{n+1}, 2^{n+1})$	$\rho^{2^n-n-2}P^{2^{n-1}k}v_n$	$n+1$

6. η-INVERTED HOMOTOPY GROUPS

From the h_1-inverted Adams E_∞-page, it is a short step to the η-inverted stable homotopy ring. First we must choose generators.

Definition 6.1. For $k \geq 0$ and $n \geq 2$, let $P^{2^{n-1}k}\lambda_n$ be an element of the Milnor-Witt $(2^{n+1}k + 2^n - 1)$-stem that is detected by $\rho^{2^n-n-2}P^{2^{n-1}k}v_n$.

There are choices in these definitions, which are measured by Adams E_∞-page elements in higher filtration. For example, there are four possible choices for λ_2 because of the presence of ρv_2 and $\rho^2 v_2$ in higher filtration.

Theorem 6.2. *As a $\mathbb{Z}_2[\eta^{\pm 1}]$-module, the η-inverted $\mathbb{R}$-motivic stable homotopy ring is generated by 1 and $P^{2^{n-1}k}\lambda_n$ for $n \geq 2$ and $k \geq 0$. The generator $P^{2^{n-1}k}\lambda_n$ lies in the Milnor-Witt $(2^{n+1}k + 2^n - 1)$-stem and is annihilated by 2^{n+1}. All products are zero, except for those involving 2 or η.*

Proof. In the η-inverted stable homotopy ring, ρ and 2 differ by a unit because $\rho\eta^2 = -2\eta$ [M2]. Therefore, the ρ-torsion information given in Proposition 5.9 translates to 2-torsion information in homotopy.

Except for 1, all $\mathbb{Z}_2[\eta^{\pm 1}]$-module generators lie in Milnor-Witt stems that are congruent to 3 modulo 4. Therefore, such generators must multiply to zero. ∎

Table 5 lists all generators through the Milnor-Witt 63-stem. The table also identifies Toda brackets that contain each generator. These Toda brackets are computed in Section 7.

Table 5 also reveals a pattern that matches the classical image of J.

Corollary 6.3. *If $k > 1$, then the η-inverted Milnor-Witt $(4k-1)$-stem is isomorphic to $\mathbb{Z}/2^{u+1}[\eta^{\pm 1}]$ as a module over $\mathbb{Z}_2[\eta^{\pm 1}]$, where u is the 2-adic valuation of $4k$.*

TABLE 5. $\mathbb{Z}_2[\eta^{\pm 1}]$-module generators for $\hat{\pi}^{\mathbb{R}}_{*,*}[\eta^{-1}]$

mw	E_∞	$\hat{\pi}^{\mathbb{R}}_{*,*}[\eta^{-1}]$	2^k-torsion	bracket	indeterminacy
0	1	1	∞		
3	v_2	λ_2	3		
7	$\rho^3 v_3$	λ_3	4	$\langle 2^3, \lambda_2, \lambda_2\rangle$	$2^3\lambda_3$
11	$P^2 v_2$	$P^2\lambda_2$	3	$\langle 2^4, \lambda_3, \lambda_2\rangle$	
15	$\rho^{10} v_4$	λ_4	5	$\langle 2^3, \lambda_2, P^2\lambda_2\rangle$	$2^3\lambda_4$
19	$P^4 v_2$	$P^4\lambda_2$	3	$\langle 2^5, \lambda_4, \lambda_2\rangle$	
23	$\rho^3 P^4 v_3$	$P^4\lambda_3$	4	$\langle 2^5, \lambda_4, \lambda_3\rangle$	
27	$P^6 v_2$	$P^6\lambda_2$	3	$\langle 2^5, \lambda_4, P^2\lambda_2\rangle$	
31	$\rho^{25} v_5$	λ_5	6	$\langle 2^3, \lambda_2, P^6\lambda_2\rangle$	$2^3\lambda_5$
35	$P^8 v_2$	$P^8\lambda_2$	3	$\langle 2^6, \lambda_5, \lambda_2\rangle$	
39	$\rho^3 P^8 v_3$	$P^8\lambda_3$	4	$\langle 2^6, \lambda_5, \lambda_3\rangle$	
43	$P^{10} v_2$	$P^{10}\lambda_2$	3	$\langle 2^6, \lambda_5, P^2\lambda_2\rangle$	
47	$\rho^{10} P^8 v_4$	$P^8\lambda_4$	5	$\langle 2^6, \lambda_5, \lambda_4\rangle$	
51	$P^{12} v_2$	$P^{12}\lambda_2$	3	$\langle 2^6, \lambda_5, P^4\lambda_2\rangle$	
55	$\rho^3 P^{12} v_3$	$P^{12}\lambda_3$	4	$\langle 2^6, \lambda_5, P^4\lambda_3\rangle$	
59	$P^{14} v_2$	$P^{14}\lambda_2$	3	$\langle 2^6, \lambda_5, P^6\lambda_2\rangle$	
63	$\rho^{56} v_6$	λ_6	7	$\langle 2^3, \lambda_2, P^{14}\lambda_2\rangle$	$2^3\lambda_6$

7. TODA BRACKETS

Even though its primary multiplicative structure is uninteresting, the η-inverted $\mathbb{R}$-motivic stable homotopy ring has rich higher structure in the form of Toda brackets. We will explore some of the 3-fold Toda bracket structure. In particular, we will show that all of the generators can be inductively constructed via Toda brackets, starting from just 2 and λ_2. Table 5 lists one possible Toda bracket decomposition for each generator up to the Milnor-Witt 63-stem.

We observed in the proof of Theorem 6.2 that the element ρ of the Adams E_∞-page detects the element 2 of the η-inverted stable homotopy ring. We will use this fact frequently in the following results.

Lemma 7.1. *The Toda bracket $\langle 2^3, \lambda_2, \lambda_2\rangle$ contains an element detected by $\rho^3 v_3$ in the Milnor-Witt 7-stem, and its indeterminacy is detected by $\rho^6 v_3$.*

Proof. Moss's convergence theorem [M3, Theorem 1.2] and the differential $d_2(v_3) = v_2^2$ show that $\langle 2^3, \lambda_2, \lambda_2\rangle$ is detected by $\rho^3 v_3$.

The indeterminacy follows from the facts that there are no multiples of λ_2 and that there is a unique multiple of 2^3 in the Milnor-Witt 7-stem. ∎

Remark 7.2. The proof of Lemma 7.1 applies just as well to show that $\langle 2^4, \lambda_3, \lambda_3\rangle$ is detected by $\rho^{10} v_4$ in the Milnor-Witt 15-stem. In higher stems, the analogous brackets do not produce generators. For example, the Massey product $\langle \rho^5, \rho^{10} v_4, \rho^{10} v_4\rangle$ is already defined in Ext, which implies that the corresponding Toda bracket must be detected in filtration least 27. However, $\rho^{25} v_5$ detects the generator of the Milnor-Witt 31-stem, and it lies in filtration 26.

Lemma 7.3. *For $n \geq 2$, the Toda bracket $\langle 2^3, \lambda_2, P^{2^{n-1}-2}\lambda_2\rangle$ is detected by $\rho^{2^{n+1}-n-3} v_{n+1}$. The indeterminacy in this Toda bracket is generated by $2^3\lambda_{n+1}$.*

Proof. This follows from Moss's convergence theorem [M3, Theorem 1.2], together with the Adams differential $d_n(\rho^{2^{n+1}-n-6}v_{n+1}) = P^{2^{n-1}-2}v_2^2$. ∎

Lemma 7.4. *For $n \geq 2$, the Toda bracket $\langle 2^{n+2}, \lambda_{n+1}, P^{2^{n-1}-2}\lambda_2\rangle$ is detected by $P^{2^n-2}v_2$. The Toda bracket has no indeterminacy.*

Proof. Lemma 4.9 implies that there is a Massey product

$$P^{2^n-2}v_2 = \langle \rho^{n+2}, \rho^{2^{n+1}-n-3}v_{n+1}, P^{2^{n-1}-2}v_2\rangle,$$

with no indeterminacy. Moss's convergence theorem [M3, Theorem 1.2] establishes the desired result. ∎

Lemma 7.5. *If $m > n \geq 2$, then the Toda bracket $\langle 2^{m+1}, \lambda_m, P^{2^{n-1}k}\lambda_n\rangle$ is detected by $\rho^{2^n-n-2}P^{2^{m-2}+2^{n-1}k}v_n$. The Toda bracket has no indeterminacy.*

Proof. Lemma 4.9 implies that there is a Massey product

$$\rho^{2^n-n-2}P^{2^{m-2}+2^{n-1}k}v_n = \langle \rho^{m+1}, \rho^{2^m-m-2}v_m, \rho^{2^n-n-2}P^{2^{n-1}k}v_n\rangle.$$

Moss's convergence theorem [M3, Theorem 1.2] establishes the desired result. ∎

Proposition 7.6. *Every generator $P^{2^{n-1}k}\lambda_n$ of the η-inverted $\mathbb{R}$-motivic stable homotopy ring can be constructed via iterated 3-fold Toda brackets starting from 2 and λ_2.*

Proof. Lemmas 7.3 and 7.4 alternately show that the generators λ_n and the generators $P^{2^n-2}\lambda_2$ can be constructed via iterated 3-fold Toda bracket starting from 2 and λ_2. Then Lemma 7.5 shows that any $P^{2^{n-1}k}\lambda_n$ can be constructed. ∎

Example 7.7. Suppose we wish to find a Toda bracket decomposition for $P^{40}\lambda_3$. Since $40 = 2^{7-2} + 2^{3-1}\cdot 4$, we can apply Lemma 7.5 with $m = 7$, $n = 3$, and $k = 4$ to conclude that $P^{40}\lambda_3$ is detected by the Toda bracket $\langle 2^8, \lambda_7, P^8\lambda_3\rangle$.

FIGURE 2. The η-inverted $\mathbb{R}$-motivic Adams E_3-page



FIGURE 3. The η-inverted $\mathbb{R}$-motivic Adams E_∞-page



References

[A] J. F. Adams, *A finiteness theorem in homological algebra*, Proc. Cambridge Philos. Soc. **57** (1961) 31–36.

[AM] M. Andrews and H. Miller, *Inverting the Hopf map in the Adams-Novikov spectral sequence*, preprint, 2014.

[DI] D. Dugger and D. C. Isaksen, *Low dimensional Milnor-Witt stems over* $\mathbb{R}$, arXiv:1502.01007, preprint, 2015.

[GI] B. Guillou and D. C. Isaksen, *The η-local motivic sphere*, J. Pure Appl. Algebra **219** (2015) 4728–4756.

[H] M. A. Hill, *Ext and the motivic Steenrod algebra over* $\mathbb{R}$, J. Pure Appl. Algebra **215** (2011) 715–727.

[I] D. C. Isaksen, *Stable stems*, arXiv:1407.8418, preprint, 2014.

[M1] J. Peter May, *Matric Massey products*, J. Algebra **12** (1969) 533–568.

[M2] F. Morel, *On the motivic π_0 of the sphere spectrum*, Axiomatic, enriched and motivic homotopy theory, NATO Sci. Ser. II Math. Phys. Chem. **131** (2004) 219–260, Kluwer Acad. Publ., Dordrecht.

[M3] R. Michael F. Moss, *Secondary compositions and the Adams spectral sequence*, Math Z. **115** (1970) 283–301.

Department of Mathematics, University of Kentucky, Lexington, KY 40506, USA
E-mail address: bertguillou@uky.edu

Department of Mathematics, Wayne State University, Detroit, MI 48202, USA
E-mail address: isaksen@wayne.edu